

AD-A150 595 ORGANOPHOSPHAZENES 18 FRIEDEL-CRAFTS PHENYLATION
REACTIONS OF ALKYL AND N. (U) VERMONT UNIV BURLINGTON
DEPT OF CHEMISTRY C W ALLEN ET AL. 15 NOV 84 TR-16
UNCLASSIFIED N00014-77-C-0605 F/G 7/3


ORGANOPHOSPHAZENES 18 FRIEDEL-CRAFTS PHENYLATION
REACTIONS OF ALKYL AND N. (U) VERMONT UNIV BURLINGTON
DEPT OF CHEMISTRY C W ALLEN ET AL. 15 NOV 84 TR-16
N00014-77-C-0605 F/G 7/3

174

NL

END

5710



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A150 595

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 16	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Organophosphazenes, 18. Friedel-Crafts Phenylation Reactions of Alkyl and Nimethylamino Fluorocyclotriphosphazenes		5. TYPE OF REPORT & PERIOD COVERED Technical Report
6. AUTHOR(s) Christopher W. Allen, Scott Bedell, William T. Pennington and A. Wallace Condes		6. PERFORMING ORG. REPORT NUMBER
7. PERFORMING ORGANIZATION NAME AND ADDRESS Departments of Chemistry, The University of Vermont, Burlington, VT 05405 and The University of Arkansas, Fayetteville, Arkansas 72701		8. CONTRACT OR GRANT NUMBER(s) N001477C-0605
8. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, VA 22217		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
9. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		10. REPORT DATE 11/15/84
		11. NUMBER OF PAGES 30
		12. SECURITY CLASS. (of this report) Unclassified
		13. DECLASSIFICATION/DOWNGRADING SCHEDULE
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale; its distribution is unlimited		
15. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
16. SUPPLEMENTARY NOTES Accepted for publication in Inorganic Chemistry		
17. KEY WORDS (Continue on reverse side if necessary and identify by block number) cyclophosphazene Friedel-Crafts Reactions nmr x-ray structure		
18. ABSTRACT (Continue on reverse side if necessary and identify by block number) The monodimethylamino and monobutylpentafluorocyclotriphosphazenes, $N_3P_3F_5R$ ($R = N(CH_3)_2$, $n-C_4H_9$, $t-C_4H_9$) undergo the Friedel-Crafts phenylation reaction to yield $N_3P_3F_4(C_6H_5)R$. A geminal configuration was assigned to each of the phenyl derivatives based on the NMR (1H , ^{13}C , ^{31}P) spectroscopic data. The crystal and molecular structures of $N_3P_3F_4(C_6H_5)(t-C_4H_9)$ have been deter- mined; the crystals are orthorhombic, space group P_{nma} , with $a = 7.264(1)\text{\AA}$.		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-660185 02 04 018
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

leg R = sub w CuA leg P10
 20. Abstract (Continued)

b = 12.204(2) Å, c = 17.028(3) Å, V = 1509.5(7) Å³, and Z = 4. The final refinement gave R = 0.042 and R_w = 0.054 for 1087 observed reflections. The molecule, which has a crystallographically-imposed mirror plane of symmetry, has the predicted geminal arrangement of organic groups and P-N distances of 1.618(1), 1.527(2), and 1.565(1) Å. The central P-N ring is planar within 0.007 Å. Bond angles are: C-P-C, 108.9(1)°; F-P-F, 96.36(9)°; N-P-N, 114.1(1); 120.91(9)°; P-N-P, 122.58(9); 118.9(1)°. The observation of Friedel-Crafts phenylation of the alkylphosphazenes demonstrates that π donation from an exocyclic substituent is not a necessary prerequisite for this reaction.

Originator furnished keywords in closure.

100-30730-114

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
Distribution/	
Availability Codes	
Avail and/or	
Special	
A-1	



OFFICE OF NAVAL RESEARCH
Contract N001477C-0605
Project NR 356-663
Technical Report No. 16

Organophosphazenes. 18. Friedel-Crafts Phenylation Reactions
of Alkyl and Dimethylamino Fluorocyclotriphosphazenes

by

Christopher W. Allen, Scott Bedell, William T. Pennington and A. Wallace Condes

Accepted for Publication in
Inorganic Chemistry

University of Vermont
Department of Chemistry
Burlington, Vermont 05405

November 15, 1984

Reproduction in whole or in part is permitted for any purposes of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

Contribution from the Departments of Chemistry,
The University of Vermont, Burlington, Vermont 05405
and
The University of Arkansas, Fayetteville, Arkansas 72701

Organophosphazenes. 18. Friedel-Crafts Phenylation Reactions of Alkyl and
Dimethylamino Fluorocyclotriphosphazenes.¹

Christopher W. Allen^{*2a}, Scott Bedell^{2a}, William T. Pennington^{2b} and
A. Wallace Cordes^{2b}

Received


Abstract

The monodimethylamino and monobutylpentafluorocyclotriphosphazenes, $N_3P_3F_5R$ ($R = N(CH_3)_2$, $n-C_4H_9$, $t-C_4H_9$) undergo the Friedel-Crafts phenylation reaction to yield $N_3P_3F_4(C_6H_5)R$. A geminal configuration was assigned to each of the phenyl derivatives based on the NMR (1H , ^{13}C , ^{31}P) spectroscopic data. The crystal and molecular structures of $N_3P_3F_4(C_6H_5)(t-C_4H_9)$ have been determined; the crystals are orthorhombic, space group P_{nma} , with $a = 7.264(1)\text{\AA}$, $b = 12.204(2)\text{\AA}$, $c = 17.028(3)\text{\AA}$, $V = 1509.5(7)\text{\AA}^3$, and $Z = 4$. The final refinement gave $R = 0.042$ and $R_w = 0.054$ for 1087 observed reflections. The molecule, which has a crystallographically-imposed mirror plane of symmetry, has the predicted geminal arrangement of organic groups and P-N distances of 1.618(1), 1.527(2), and 1.565(1) \AA . The central P-N ring is planar within 0.007 \AA . Bond angles are: C-P-C, 108.9(1) $^\circ$; F-P-F, 96.36(9) $^\circ$; N-P-N, 114.1(1); 120.91(9) $^\circ$; P-N-P, 122.58(9); 118.9(1) $^\circ$. The observation of Friedel-Crafts phenylation of the alkylphosphazenes demonstrates that π donation from an exocyclic substituent is not a necessary prerequisite for this reaction.

Introduction

The Friedel-Crafts arylation is a classic reaction³ in phosphazene chemistry dating back to the pioneering studies of Bode and Bach.⁴ This reaction can be used to produce geminal aryl groups in chloro^{4,5} and fluorocyclophosphazenes.⁶ The observation that the reaction usually yields geminal di, tetra and hexaaryl phosphazenes but not mono, tri or penta substituted derivatives leads to the suggestion that a $\equiv \text{PPhCl}$ center is more reactive than a $\equiv \text{PCl}_2$ center. Aminochlorophosphazenes also undergo the Friedel-Crafts arylation reaction which allows for conversion of a $\equiv \text{PClNR}_2$ center to a $\equiv \text{P(Ph)NR}_2$ center.⁷ Depending on the nature of the amine, side reactions may also become significant. The phosphazo substituent also promotes Friedel-Crafts arylation at a $\equiv \text{PCl(N=PPh}_3\text{)}$ center.⁸

In spite of the synthetic utility of this reaction, little is known about the mechanism. The first step of the reaction is often assumed to be the formation of a phosphonium ion arising from halide abstraction by the Lewis acid. Definitive evidence on this point has yet to become available. The



addition compounds of aluminum tribromide with halocyclophosphazenes, $\text{N}_3\text{P}_3\text{X}_6 \cdot n\text{AlBr}_3$ (X = Cl, n = 1; X = Br, n = 1,2) are believed to have the aluminum tribromide entity coordinated to an endocyclic nitrogen atom.⁹

Related addition compounds of SbCl_5 and TiCl_4 apparently involve the phosphonium ion.¹⁰ Once the phosphonium ion forms attack on the solvent (benzene) could lead to a cationic sigma complex and deprotonation to the observed arylphosphazene. The assumption is implicitly or explicitly made that the facility of the Friedel-Crafts reaction at a $\equiv \text{PXR}$ (R = aryl, amino; X = F, Cl) center is due to stabilization of the positive charge (or

developing positive charge) on the phosphorus atom by the electron donating substituent, R.

In this paper, we wish to explore the nature of some of the electronic factors involved in the proposed stabilization of the positive charge on the phosphorus center and expand the range of systems which undergo the Friedel-Crafts arylation reactions.

Experimental Section

Materials and Methods. Hexachlorocyclotriphosphazene, $N_3P_3Cl_6$ (Firestone Corp.), was converted to hexafluorocyclotriphosphazene¹¹, which was in turn converted to (dimethylamino)pentafluorocyclotriphosphazene, $N_3P_3F_5N(CH_3)_2$ ¹², (n-butyl)pentafluorocyclotriphosphazene, $N_3P_3F_5CH_2(CH_2)_2CH_3$ ¹³, and (t-butyl)pentafluorocyclotriphosphazene, $N_3P_3F_5C(CH_3)_3$ ¹³, by previously reported procedures. Diethylether was distilled from sodium/benzophenone. Benzene¹⁴ was distilled from sodium. Other reagents and solvents were obtained from standard sources and used without further purification. Nmr spectra (in $CDCl_3$) were reported on a Bruker WM250 spectrometer operating at 250.1 (1H), 62.9 (^{13}C), 235.2 (^{19}F), and 101.2 MHz (^{31}P). Tetramethylsilane, Me_4Si (for 1H and ^{13}C), and fluorotrichloromethane, $CFCl_3$ (for ^{19}F), were used as internal references. For ^{31}P nmr, 85% H_3PO_4 was used as an external reference. Chemical shifts upfield to the reference are assigned a negative sign. ^{13}C , ^{19}F , and ^{31}P nmr spectra were recorded under conditions of broad-band decoupling. Mass spectra were recorded on a Finnegan 4610 mass spectrometer operating at 80eV. Elemental analyses were performed by Integral Microanalytical Laboratories.

Preparations of $N_3P_3F_5[N(CH_3)_2]C_6H_5$ (1). A solution of 2.65g (.01 mol) of $N_3P_3F_5N(CH_3)_2$ in 12 mL of benzene was added to a solution of triethylamine (2.8g, 0.03 mol) and anhydrous aluminum chloride (9.3g, 0.07 mol) in 50 mL of benzene.¹⁴ The reaction mixture was allowed to stir at reflux for five days. The mixture was hydrolyzed over acidified ice water and the layers were separated. The benzene layer was washed sequentially with saturated sodium bicarbonate solution and distilled water then dried over anhydrous sodium

sulfate and decolorized. After removal of the solvent, the yellow liquid was distilled at reduced pressure (0.02 mmHg) to yield 1.0g (31.3% of theory) of a colorless oil, bp 60°C (0.02 mmHg). Anal. Calcd. for $N_4P_3F_4C_8H_{11}$: C, 28.92; H, 3.31; mol wt 332. Found: C, 28.93; H, 3.49; mol wt 332 (mass spectrum¹⁵). 1H NMR: $^{16}\delta_{N(CH_3)_2} = 2.6$ (d, 6H), $^3J_{PH} = 12.9$; $\delta_{C_6H_5-m,p} = 7.5$ (m, 3H); $\delta_{C_6H_5-o} = 7.8$ (q, 2H), $^3J_{PH} = 13.8$. ^{13}C NMR: $\delta_{N(CH_3)_2} = 35.6$ (d), $^2J_{PC} = 3.5$; $\delta_{ArC_2} = 131.2$ (d); $^2J_{PC} = 10.7$; $\delta_{ArC_3} = 128.8$ (d), $^3J_{PC} = 14.9$; $\delta_{ArC_4} = 132.7$ (d), $^4J_{PC} = 3.1$. ^{31}P NMR: $\delta_{\equiv PF_2} = 10.2$ (m, 2P), $^1J_{PF} = 916.0$ ¹⁷; $\delta_{\equiv PN(CH_3)_2C_6H_5} = 29.5$ (m, 1P), $^2J_{pp} = 63.7$.

Preparation of $N_3P_3F_4[CH_2(CH_2)_2CH_3]C_6H_5$ (2). The procedure was identical with that described for the preparation of **1** except that $N_3P_3F_5CH_2(CH_2)_2CH_3$ was used in place of $N_3P_3F_5N(CH_3)_2$. In a typical experiment, the following quantities were allowed to react for five days: $N_3P_3F_5CH_2(CH_2)_2CH_3$ (2.00g, 0.007 mol), triethylamine (2.25g, .02 mol) and aluminum chloride (7.45g, .06 mol). Distillation yielded a colorless oil whose ^{31}P nmr spectrum consisting of largely the unreacted starting material, small amounts of $N_3P_3F_4[CH_2(CH_2)_2CH_3]C_6H_5$ (less than 10%) and traces of other unidentified materials. Careful fractionation (35° at 0.02 mmHg) yielded an impure sample of the desired product. Anal. Calcd. for $N_3P_3F_4C_{10}H_{14}$: C, 34.78; H, 4.06; mol wt 345. Found: C, 29.43; H, 4.50; mol wt 345 (mass spectrum¹⁵). 1H NMR: $^{16}\delta_{CH_2(CH_2)_2CH_3} = 1.9$ (m, 2H); $\delta_{CH_2CH_2CH_2CH_3} = 1.5$ (m, 4H); $\delta_{(CH_2)_3CH_3} = 0.9$ (m, 3H); $\delta_{C_6H_5-m,p} = 7.5$ (m, 3H); $\delta_{C_6H_5-o} = 7.8$ (q, 2H), $^3J_{PH} = 13.3$. ^{13}C NMR: $\delta_{C\alpha} = 33.6$ (m), $^1J_{PC} = 98.1$; $\delta_{C_{\beta,\gamma}} = 23.2$ (m, 2C), $\delta_{C_{\Delta}} = 13.5$ (s); $\delta_{ArC_2} = 130.0$ (d), $^2J_{PC} = 11.0$; $\delta_{ArC_3} = 128.8$ (d), $^3J_{PC} = 12.9$; $\delta_{ArC_4} = 132.6$ (d), $^4J_{PC} = 3.4$. ^{31}P NMR (approximate): $\delta_{\equiv PF_2} = 7.1$, $^1J_{PF} = 883$ ¹⁷; $\delta_{\equiv P(C_4H_9)C_6H_5} = 37.8$, $^2J_{pp} = 36$.

Preparation of $N_3P_3F_4[C(CH_3)_3]C_6H_5$ (3). The procedure was identical with that described above except that $N_3P_3F_5C(CH_3)_3$ was used as the phosphazene starting material. In a typical experiment, the following quantities were allowed to react for five days: $N_3P_3F_5C(CH_3)_2$ (3.80g, 0.013 mol), triethylamine (4.10g, 0.04 mol), and aluminum chloride

(13.5g, 0.10 mol). The yellow solid remaining after removal of the benzene was sublimed to yield 1.78g (39.0% of theory) of a milk-white solid, mp 102-103°. Anal. Calcd. for $N_3P_3F_4C_{10}H_{14}$: C, 34.78; H, 4.06; mol wt 345. Found: C, 35.05; H, 3.68; mol wt 345 (mass spectrum¹⁵).

1H NMR¹⁶: $\delta_{C(CH_3)_3}$ 1.1 (d, 9H), $^3J_{PH} = 17.7$; $\delta_{C_6H_5-m,p} = 7.5$ (m, 3H); $\delta_{C_6H_5-o} = 7.8$ (q, 2H), $^3J_{PH} = 12.0$. ^{13}C NMR: $\delta_{C(CH_3)_3} = 33.9$ (m); $\delta_{ArC_2} = 131.9$ (d), $^2J_{PC} = 10.1$; $\delta_{ArC_3} = 128.2$ (d), $^3J_{PC} = 12.9$; $\delta_{ArC_4} = 132.5$ (d), $^4J_{PC} = 3.4$. ^{31}P NMR: $\delta_{PF_2} = 9.2$, $^1J_{PF} = 885.1$ ¹⁷; $\delta_{PPhC(CH_3)_3} = 46.1$, $^3J_{PP} = 36.1$.

Attempted Friedel-Crafts reaction of $N_3P_3F_5OCH_2CH_3$. The procedure was identical with that described above except that $N_3P_3F_5OCH_2CH_3$ (prepared from $N_3P_3F_6$ and $LiOCH_2CH_3$) was used as the phosphazene starting material. No phosphazene containing products were recovered from the benzene layer after hydrolysis.

X-ray Analysis of $N_3P_3F_4(C_6H_5)(t-C_4H_9)$. Crystals of 3 suitable for x-ray work were obtained by recrystallization from a petroleum ether- CH_2Cl_2 mixture. A colorless crystal of approximate dimensions 0.30 X 0.30 X 0.35 mm, mounted on a glass fiber, was used for data collection on an Enraf-Nonius CAD-4 diffractometer employing graphite-monochromated $Mo K\alpha$ ($\gamma = 0.71073 \text{ \AA}$) radiation. A least-squares fit of the diffractometer setting angles for 25 carefully centered reflections ($24^\circ < 2\theta < 37^\circ$) gave the unit cell parameters listed in Table I.

Systematic absences of $0kl$ ($k+1$, odd) and $hk0$ (h odd) indicate the space group $Pna2_1$ (No. 33) or $Pnma$ (No. 62); the centrosymmetric space group $Pnma$ was verified by the refinement. Reflections were measured using ω - 2θ scans for 2θ from 2 to 50° ($h = 0$ to 8, $k = 14$ to 0, $l = 0$ to 20). The scan range was $(1.00 + 0.35 \tan \theta)^\circ$ and the scan speed varied from 20 to $4^\circ/\text{min}$. Of the 1529 reflections measured, 1087 had $I > 3\sigma(I)$ and were used in the refinement. Data were corrected for Lorentz and polarization effects, but due to the low linear absorption coefficient and regular shape of the crystal an absorption correction was not made. Periodic measurement of the intensities of three reflections (2 9 0, 4 0 4, 0 5 11) indicated a 28% decline during the 13.2 hours of data collection, and intensities were

corrected accordingly.

The structure was solved by direct methods (MULTAN80)¹⁸ and refined by full-matrix least-squares techniques. The molecule has a crystallographically imposed mirror symmetry. Hydrogen atoms were located using difference Fourier techniques and were included in the refinement. The final cycle of refinement based on $(|F_o| - |F_c|)^2$ included positional and anisotropic thermal parameters for all non-hydrogen atoms and positional parameters for the hydrogen atoms. The isotropic thermal parameters of each hydrogen atom was constrained at a value 20% greater than the B_{eq} of the carbon atom to which it was attached. The final $R^{19} = 0.042$ and $R_w^{20} = 0.054$. The weighting scheme used was $w = [\sigma^2(F) + 0.03F^2]^{-1}$ where $\sigma(F)$ was derived from counting statistics. The goodness of fit²¹ was 2.36. In the final cycle of refinement the maximum shift/error was 0.09. The final difference map has a maximum value of $0.36 \text{ e}^-/\text{\AA}^3$. No secondary extinction correction was made. The atomic scattering factors for neutral atoms were those of Cromer and Waber²² and the real and imaginary dispersion corrections were those of Cromer.²³ The computer programs used were those provided by the Enraf-Nonius Structure Determination Package.

Results and Discussion

The synthetic results show that a fluorine atom in both monoalkyl and mono(N,N-dimethylamino)pentafluorocyclotriphosphazenes can be conveniently replaced by a phenyl group under Friedel-Crafts conditions. Although dialkylaminochlorophosphazenes have previously been shown to undergo Friedel-Crafts phenylation⁷, we report the first example of the corresponding reaction in the fluorophosphazene series and the first example of Friedel-Crafts phenylation of an alkylphosphazene. The new compounds (**1**, **2**, **3**) have all been characterized by elemental analyses, mass spectrometry and nmr (¹H, ¹³C, ³¹P) spectroscopy. Observations of the stereochemical course of the Friedel-Crafts phenylation of other cyclophosphazenes⁴⁻⁷ leads to the suggestion of a geminal substitution pattern being followed in the synthesis of **1**, **2** and **3**. The observed values of the $^2J_{\text{PCC}}$ ²⁴ and $^3J_{\text{PCH}}$ ²⁵ coupling constants in **1**, **2** and **3** are consistent

with geminal rather than non-geminal configurations. The ^{31}P nmr spectra of each of the new compounds shows a large, complex, triplet ($J_{\text{PF}} \sim 883 - 916 \text{ Hz}$) due to the $\equiv\text{PF}_2$ centers and a small ($J_{\text{PP}} \sim 36 \text{ Hz}$) triplet due to the $\equiv\text{PR}(\text{Ph})$ center. The lack of a doublet due to a $\equiv\text{PFR}$ (or $\equiv\text{PFPh}$) center confirms the absence of non-geminal products.

The crystal and molecular structures of **3** were determined in order to verify the proposed geminal configuration and to explore the structural consequences of the sterically crowded environment arising from substitution of both a phenyl and tert-butyl group on the same phosphorus atom. The final atomic coordinates of all unique atoms are given in Table II. Selected bond lengths and bond angles may be found in Table III and an ORTEP drawing of the molecule, together with the atom numbering scheme, is shown in Figure 1. The observed molecular structure of **3** confirms the geminal configuration of organic functions on the phosphazene. The solid state structure of **3** is such that the phosphazene ring is bisected by a mirror plane containing N_1 , P_1 , C_1 , C_4 , H_4 , C_5 , C_7 and H_{71} . The conformations adopted by the phenyl and tert-butyl groups lead to the minimum amount of steric interaction between the two groups. A comparison of the structure of **3** with the structure of the previously reported geminal diphenyltetrafluorocyclotriphosphazene, $2,2\text{-N}_3\text{P}_3\text{F}_4(\text{C}_6\text{H}_5)_2$ (**4**)²⁶ is of interest. In **4** five of the six phosphazene ring atoms are close to coplanarity with the phenyl substituted phosphorus atom 0.20\AA out of this plane. In **3** the central ring is more nearly planar (within 0.007\AA): the P_3N_2 segment which excludes the unique nitrogen atom is planar within 0.001\AA and the $\text{N } 1$ atom is displaced only $0.014(3)\text{\AA}$ from this plane. In comparing **3** to **4** a small effect of the increased steric demand of the tert-butyl group over the phenyl group is seen in the bond angles at the organosubstituted phosphorus atom ($\angle\text{CPC} = 108.9(1)$ for **3** and $107.9(3)$ for **4**; $\angle\text{NPN} = 114.1(1)$ for **3** and

115.3 (3) for 4). The phosphorus-nitrogen bond lengths are virtually identical in 3 and 4 and vary in a manner which is typical of assymmetrically substituted phosphazenes and can be correlated with the relative ability of each phosphorus center to attract nitrogen lone pair electron density.²⁶ The small differences in P-N distances of P2 are consistent with the effect of replacing an sp^2 carbon atom (in 4) by a sp^3 carbon atom (in 3) resulting in a decrease of electron attracting power of the organosubstituted phosphorus atom. Consequently, the lone pair of electrons at N(2) is more strongly attracted to P(2) in 3 than in 4. The increased charge density on P(2) in 3 results in less charge delocalization from N(1) and hence a longer P(2)N(1) distance in 3 vs 4 (1.565(1) vs 1.555(4)).

One of the goals of this study was to probe substituent electronic effects which may be operative in the Friedel-Crafts phenylation reaction. Since previous investigations have employed exocyclic groups which are potential π donors to the phosphorus center, monoalkylpentafluorocyclotriphosphazenes were examined in order to clarify differences between σ and π donor substituent effects. The fact that the alkylphosphazenes undergo the phenylation reaction unambiguously demonstrates that π donation from the exocyclic function is not a necessary prerequisite for this reaction to be effective. The significantly higher yield in the reaction of tert-butyl versus the n-butyl phosphazene suggests the operation of a mechanism wherein the phosphorus atom undergoing substitution goes to a lower coordinate intermediate (or transition state). This process would be most favorable for the tert-butyl derivative where the maximum relief of steric strain can be obtained. A reasonable model for the Friedel-Crafts phenylation reaction of substituted phosphazenes is one in which the substituted phosphorus atom goes through a three coordinate phosphorus (v) intermediate which is stabilized by the σ electron releasing nature of the substituent. It is possible that in the phenyl and dialkylamino systems, the σ electron releasing effect may be supplemented by a π effect but conclusive evidence of this has yet to be presented. The reluctance of the

monoethoxypentafluorocyclotriphosphazene to enter into the Friedel-Crafts reaction is consistent with these proposals since the high electronegativity of the oxygen atom makes it a poor electron releasing substituent and the small size does not allow for significant relief of steric strain on going to a lower coordinate transition state intermediate.

Acknowledgements.

Phosphazene chemistry at the University of Vermont is supported in part by the Office of Naval Research. We wish to thank Firestone Corporation for a generous gift of hexachlorocyclotriphosphazene and Mr. Weglarz for checking certain synthetic results. The X-ray work at Arkansas was supported by the State of Arkansas and the National Science Foundation (EPSCOR grant ISP801147).

Supplementary Material Available. Table 5.1. showing major mass spectral fragments and their intensities (pages), Table 5.2 showing structure factor amplitudes for $N_3P_3F_4(C_6H_5)(C_4H_9)$, and Table 5.3 showing anisotropic thermal parameters for $N_3P_3F_4(C_6H_5)(C_4H_9)$. Ordering information is given on any current masthead page.

References

1. Part 17. Allen, C.W., Descorcie, J. L., Ramachandran, K.
J. Chem. Soc., Dalton Trans., in press.
2. (a) University of Vermont. (b) University of Arkansas
3. Allcock, H. R. "Phosphorus-Nitrogen Compounds"; Academic Press: New York, 1972; Krishnamurthy, S. S., Sau, A. C.; Woods, M. Adv. Inorg. Chem. Radiochem. 1978, 21, 41.
4. Bode, H.; Bach, H. Chem. Ber. 1942, 75, 215.
5. Acock, K. G.; Shaw, R. A.; Wells, F.B.G. J. Chem. Soc. 1964, 121;
Grushkin, B.; Sanchez, M.G.; Ernest, M. V.; McClanahan, J. L.; Ashby, G. E.;
Rice, R. G. Inorg. Chem. 1965, 4, 1538; McBee, E. T.; Okuhara, K.; Morton,
C. J. Inorg. Chem. 1965, 4, 1672.
6. Allen, C. W.; Tsang, F. Y.; Moeller, T. Inorg. Chem. 1968, 7, 2183; Allen,
C. W.; Moeller, T. Inorg. Syn. 1970, 12, 293; Allen, C. W.; Brunst, G. E.;
Perlman, M. E. Inorg. Chimica Acta 1980, 41, 265; Allen, C. W.; Toch, P. L.
Inorg. Chem. 1981, 20, 8.
7. Das, S.; Shaw, R. A.; Smith, B. C. Angew. Chem. Int. Ed. Engl. 1963, 7, 887;
Das, S.; Shaw, R. A.; Smith, B. C. J. Chem. Soc., Dalton Trans. 1981, 107;
Das, S.; Hasan, M.U.; Shaw, R. A.; Smith, B. C.; Woods, Z. Naturforsch.
1979, 346, 58.
8. Biddlestone, M.; Shaw, R. A. J. Chem. Soc., Dalton Trans. 1973, 2740.
9. Coxon, G. E.; Sowerby, D. B. J. Chem. Soc. A. 1969, 3012.
10. Kravchenko, E. A.; Levin, B. V.; Bananuanly, S. I.; Toktomatov, T. A.
Koord. Khim., 1977, 3, 374; Chem. Abs. 1977, 86, 182280.
11. Moeller, T.; John, K.; Tsant, F. Chem. Ind. (London), 1961, 347.
12. Glemser, O.; Niecke, E.; Roesky, H. W. J. Chem. Soc. D. 1969, 282.
13. Ramachandran, K.; Allen, C. W. J. Am. Chem. Soc. 1982, 104, 2396.
14. Benzene is a known carcinogen and hence should be only used in a well
ventilated hood while wearing appropriate protective gloves.

15. Mass spectral data are available as supplementary material.
16. All nmr (^1H , ^{31}P) chemical shifts in ppm and coupling constants in Hz.
17. Due to considerable second order character, the $^1J_{\text{PF}}$ values are only approximate.
18. Main, P.; Fiske, S. J.; Hull, S. E.; Lessinger, L.; Germain, G.; Declercq, J. P.; and Woolfson, M. M. 1980 MULTAN80- a system of computer programs for the automatic solution of crystal structures from X-ray diffraction data. University of York, England and University of Louvain, Belgium.
19. $R = \Sigma(|F_o| - |F_c|)/\Sigma|F_o|$
20. $R_w = [\Sigma w(|F_o| - |F_c|)^2/\Sigma w|F_o|^2]^{1/2}$
21. G.O.F. = $[\Sigma w(|F_o| - |F_c|)^2/N_o - N_v]^{1/2}$, where N_o = no. of observations, and N_v = no. of variables.
22. Cromer, D. T. and Waber, J. T. International Tables for X-Ray Crystallography, Vol. IV, The Kynoch Press, Birmingham, England, 1974, Table 2. 2B.
23. Cromer, D. T. International Tables for X-Ray Crystallography, Vol. IV, The Kynoch Press, Birmingham, England, 1974, Table 2.3.1.
24. Allen, C. W. J. Organometal. Chem. 1977, 125, 215.
25. Allen, C. W.; White, A. J. Inorg. Chem. 1974, 13, 1220.
26. Allen, C. W.; Faught, J. B.; Moeller, T.; Paul, I. C. Inorg. Chem. 1969, 8, 1719.




Figure 1. ORTEP drawing (50% probability ellipsoids) of 2,2-N₃P₃F₄(C₆H₅)(C₄H₉) showing the atom numbering scheme.

Table I. Crystallographic Data

formula	$C_{10}H_{14}F_4N_3P_3$
fw	345.16
cryst. syst.	orthorhombic
space group; molecules/cell	$Pnma$; 4
a, Å	7.264 (1)
b, Å	12.204 (2)
c, Å	17.028 (3)
V, Å ³	1509.5 (7)
reflens used for unit cell; 2θ range, deg	25; 24 - 37
calcd density, g cm ⁻³	1.52
F(000)	704
reflens scanned; obsd	1529; 1087
2θ range of reflens scanned, deg	2 - 50
μ (Mo Kα), cm ⁻¹	4.2
final R; R _w	0.042; 0.054
max shift in final cycle (σ)	0.09
max peak (final diff map), eÅ ⁻³	0.36
goodness of fit	2.36
parameters refined; parameter/reflen ratio	122; 8.9

Table II. Positional Parameters

Atom	x/a	y/b	z/c	Beq
P1	0.9951(1)	0.250	0.52490(5)	3.57(2)
P2	0.8471(1)	0.13958(6)	0.65138(4)	4.48(2)
F1	0.6759(3)	0.0679(2)	0.6496(1)	7.04(4)
F2	0.9549(3)	0.0696(2)	0.7091(1)	7.84(5)
N1	0.7965(5)	0.250	0.6928(2)	5.21(8)
N2	0.9409(3)	0.1388(2)	0.5711(1)	4.32(5)
C1	0.8770(5)	0.250	0.4331(2)	3.80(8)
C2	0.8298(4)	0.1522(2)	0.3968(2)	5.11(7)
C3	0.7363(5)	0.1539(3)	0.3267(2)	7.05(9)
C4	0.6923(7)	0.250	0.2920(3)	7.4(1)
C5	1.2400(5)	0.250	0.5068(2)	4.75(9)
C6	1.2928(5)	0.1476(3)	0.4606(2)	6.62(8)
C7	1.3385(7)	0.250	0.5869(3)	9.6(2)
H2	0.854(4)	0.085(2)	0.421(2)	6*
H3	0.717(4)	0.103(3)	0.304(2)	8*
H4	0.624(8)	0.250	0.254(3)	11*
H61	1.233(4)	0.156(3)	0.412(2)	7*
H62	1.422(5)	0.150(3)	0.452(2)	7*
H63	1.251(4)	0.079(3)	0.491(2)	7*
H71	1.480(8)	0.250	0.571(4)	11*
H72	1.294(5)	0.194(3)	0.625(2)	11*

Starred atoms were refined isotropically.

Table III. Selected distances (Å) and Angles (deg) for $N_3P_3F_4(C_6H_5)(C_4H_9)^a$

P1 - N2	1.618(1)	N2 - P1 - N2	114.1(1)
P1 - C1	1.784(3)	N2 - P1 - C1	108.04(7)
P1 - C5	1.806(3)	N2 - P1 - C5	108.82(8)
P2 - F1	1.521(1)	C1 - P1 - C5	108.9(1)
P2 - F2	1.520(1)	F1 - P2 - F2	96.36(9)
P2 - N1	1.565(1)	F1 - P2 - N1	108.2(1)
P2 - N2	1.527(2)	F1 - P2 - N2	110.06(8)
C1 - C2	1.387(2)	F2 - P2 - N1	108.3(1)
C2 - C3	1.374(3)	F2 - P2 - N2	110.21(8)
C3 - C4	1.351(4)	N1 - P2 - N2	120.91(9)
C5 - C6	1.526(3)	P2 - N1 - P2	118.9(1)
C5 - C7	1.540(5)	P1 - N2 - P2	122.58(9)
C-H (ave)	0.93(9)	P1 - C1 - C2	120.6(1)
		P1 - C5 - C6	109.6(2)
		P1 - C5 - C7	107.9(2)

^aNumbers in parentheses are estimated standard deviations in the least significant digits.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		

ABSTRACTS DISTRIBUTION LIST, 356E

Professor A. G. MacDiarmid
Department of Chemistry
University of Pennsylvania
Philadelphia, Pennsylvania 19174

Dr. E. Fischer, Code 2853
Naval Ship Research and
Development Center
Annapolis, Maryland 21402

Professor H. Allcock
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Professor R. Lenz
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts 01002

Professor M. David Curtis
Department of Chemistry
University of Michigan
Ann Arbor, Michigan 48105

Dr. J. Griffith
Naval Research Laboratory
Chemistry Section, Code 6120
Washington, D.C. 20375

Professor G. Wnek
Department of Materials Science
and Engineering
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Mr. Samson Jennekke
Honeywell Corporate Technology Center
10701 Lyndale Avenue South
Bloomington, Minnesota 55420

Dr. Richard M. Laine
SRI International
333 Ravenswood Avenue
Menlo Park, California 94025

Dr. John McGrath
Department of Chemical Engineering
Virginia Polytechnic Institute
Blacksburg, Virginia 24061

Dr. Adolf Amster
Chemistry Division
Naval Weapons Center
China Lake, California 93555

Dr. William Tolles
Code 6100
Naval Research Laboratory
Washington, D.C. 20375

Professor T. Katz
Department of Chemistry
Columbia University
New York, New York 10027

Professor J. Salamone
Department of Chemistry
University of Lowell
Lowell, Massachusetts 01854

Professor J. Chien
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts 01854

Professor William R. Krigbaum
Department of Chemistry
Duke University
Durham, North Carolina 27706

Dr. R. Miller
IBM Research Laboratory K42/282
5600 Cottle Road
San Jose, California 95193

ABSTRACTS DISTRIBUTION LIST, 356B

Professor T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Professor Malcolm B. Polk
Department of Chemistry
Atlanta University
Atlanta, Georgia 30314

Dr. Kurt Baum
Fluorochem, Inc.
680 S. Ayon Avenue
Azusa, California 91702

Professor H. Ishida
Department of Macromolecular Science
Case Western University
Cleveland, Ohio 44106

Professor Stephen Wellinghoff
Department of Chemical Engineering
University of Minnesota
Minneapolis, Minnesota 55455

Professor G. Whitesides
Department of Chemistry
Harvard University
Cambridge, Massachusetts 02138

Dr. K. Paciorek
Ultrasystems, Inc.
P.O. Box 19605
Irvine, California 92715

Professor H. Hall
Department of Chemistry
University of Arizona
Tucson, Arizona 85721

Professor D. Seyferth
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

END

FILMED

3-85

DTIC